



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

types and a taxonomic presentation of the group, in which *Gnetopsis elliptica* is placed provisionally with *Conostomum* in the "Conostomeae." A brief discussion of the "pollination mechanisms" of the Lagenostomales calls attention to the three distinct types exhibited by the group: that in which the free but approximated lobes of the integument surrounded and overtopped the pollen chamber and probably at pollination formed a funnel (*Physostoma*); that in which a massive "canopy" was pierced by a long micropyle (*Conostoma*); and that in which the compact canopy closely invested the conical pollen chamber, whose orifice reached to the outer surface (*Lagenostoma*). A glossary is provided at the close of the paper, since such terms as the "blow-off layer," "lagenostome," and "plinth" are not easily separated from the well-worn terms heretofore applied to the same structures.—J. M. C.

**Anatomy of Azolla.**—QUEVA<sup>36</sup> has investigated the vascular anatomy of *Azolla filiculoides*, and has secured some interesting facts. The vascular elements are differentiated in the floating, dorsiventral stem, those of the dorsal region being tracheids of small caliber, and those of the ventral region being vessels of large caliber. The transverse section of the xylem is circular, the circle being incomplete alternately on the right and left sides in the dorsal region; so that the section is really an arc which is open alternately right and left, corresponding to the alternating leaf traces. The heavy vessels of the ventral region are connected exclusively with the roots. The interpretation suggested is that the dorsal group of vessels represents a reduced bipolar group, connected at the poles with leaf traces; and that the ventral group is merely an "apolar" mass related to the roots. The amount of vascular tissue retained would seem to be a remarkable feature in a stem with such an extremely hydrophytic habit.—J. M. C.

**Germination of Helianthus.**—MILLER<sup>37</sup> has studied the transformations of the reserve materials of the sunflower during germination. The work shows both chemical and biological excellence. Main emphasis is put upon the transformation of fats. The fats extracted from the cotyledons show low acid values at all times, while those from the hypocotyl very early, and continuously thereafter, show high acid values. It was not determined whether the fats are translocated as such or as hydrolyzed products. The iodine value of the fats falls as germination advances, due as the author believes to the absorption of oxygen. As germination progresses, the fats decrease rather rapidly, while the carbohydrates increase. This furnishes further evidence for the established view that during germination fats are transformed to carbohydrates.—WILLIAM CROCKER.

<sup>36</sup> QUEVA, C., *L'Azolla filiculoides* Lam., étude anatomique. Mém. Soc. Hist. Nat. Autun **23**: pp. 24. figs. 22. 1910.

<sup>37</sup> MILLER, EDWIN C., A physiological study of the germination of *Helianthus annuus*. Annals of Botany **24**: 693-726. 1910.